



Journal of Cellular Physiology

HDAC8 regulates canonical Wnt pathway to promote differentiation in skeletal muscles

Journal:	<i>Journal of Cellular Physiology</i>
Manuscript ID	Draft
Wiley - Manuscript type:	Original Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Ferrari, Luca; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Bragato, Cinzia; Fondazione IRCCS Istituto Neurologico C. Besta, Milano, Italy; PhD program in Neuroscience, University of Milano-Bicocca Brioschi, Loredana; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Spreafico, Marco; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Esposito, Simona; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Pezzotta, Alex; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Pizzetti, Fabrizio; Department of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Italy. Moreno Fortuny, Artal; Division of Cell Matrix Biology & Regenerative Medicine, FBMH, University of Manchester. UK.; Developmental Genetics, Department of Biomedicine, University of Basel, Basel, Switzerland. Giordano, Antonio; Temple University-Sbarro Institute for Cancer Research and Molecular Medicine, Biology Bellipanni, Gianfranco; College of Science and Technology Temple University, Sbarro Institute for Cancer Research and Molecular Medicine ; College of Science and Technology Temple University, Department of Biology Riva, Paola; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Frabetti, Flavia; Department of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Italy. Viani, Paola; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Cossu, Giulio; Division of Cell Matrix Biology & Regenerative Medicine, FBMH, University of Manchester. UK. Mora, Marina; Fondazione IRCCS Istituto Neurologico C. Besta, Milano, Italy Marozzi, Anna; Università degli Studi di Milano, Dipartimento di Biotecnologie Mediche e Medicina Traslazionale Pistocchi, Anna; University of Milan, Medical Biotechnologies and Translational Medicine; University of Milan, Biosciences</p>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Key Words:	HDAC8, zebrafish, WNT/beta catenin, skeletal muscle

SCHOLARONE™
Manuscripts

For Peer Review

HDAC8 regulates canonical Wnt pathway to promote differentiation in skeletal muscles

Luca Ferrari^{1*}, Cinzia Bragato^{2,5*}, Loredana Brioschi^{1*}, Marco Spreafico¹, Simona Esposito¹, Alex Pezzotta¹, Fabrizio Pizzetti³, Artal Moreno-Fortuny^{4,6}, Gianfranco Bellipanni^{7,8}, Antonio Giordano^{7,8}, Paola Riva¹, Flavia Frabetti³, Paola Viani¹, Giulio Cossu⁴, Marina Mora², Anna Marozzi¹, Anna Pistocchi^{1#}

¹ Dipartimento di Biotecnologie Mediche e Medicina Traslazionale, Università degli Studi di Milano, Italy.

² Fondazione IRCCS Istituto Neurologico C. Besta, Milano, Italy.

³ Department of Experimental, Diagnostic and Specialty Medicine, University of Bologna, Italy.

⁴ Division of Cell Matrix Biology & Regenerative Medicine, FBMH, University of Manchester, UK.

⁵ PhD program in Neuroscience, University of Milano-Bicocca.

⁶ Developmental Genetics, Department of Biomedicine, University of Basel, Basel, Switzerland.

⁷ Department of Biology, College of Science and Technology, Temple University, Philadelphia, Pennsylvania.

⁸ Sbarro Institute for Cancer Research and Molecular Medicine, College of Science and Technology, Temple University, Philadelphia, Pennsylvania.

*The authors equally contributed to this work

Correspondence to: anna.pistocchi@unimi.it

Running Title: HDAC8 role in skeletal muscle differentiation

Acknowledgements

We thank P.L. Lollini, University of Bologna, for providing rhabdomyosarcoma cell lines; Cotelli F. and Mazzola M., University of Milan, for the priceless advices, practical help and useful discussion of the zebrafish data. This work was supported by the AIRC, Associazione Italiana per la Ricerca sul Cancro (MFAG#18714). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

Keywords: HDAC8, skeletal muscle, rhabdomyosarcoma, Wnt, zebrafish

Abstract

Histone deacetylase 8 (HDAC8) is a class 1 histone deacetylase and a member of the cohesin complex. *HDAC8* is expressed in smooth muscles but its expression in skeletal muscle has not been described. We show for the first time that *HDAC8* is expressed in human and zebrafish skeletal muscles. Using RD/12 and RD/18 rhabdomyosarcoma cells with low and high differentiation potency respectively, we highlight a specific correlation with *HDAC8* expression and an advanced stage of muscle differentiation. We inhibit HDAC8 activity through the specific PCI-34051 inhibitor in murine C2C12 myoblasts and zebrafish embryos and we observed skeletal muscles differentiation impairment. We also found a positive regulation of the canonical Wnt signalling by HDAC8 that might explain muscle differentiation defects. These findings suggest a novel mechanism through which HDAC8 expression in a specific time window of skeletal muscle development positively regulates canonical Wnt pathway that is necessary for muscle differentiation.

1 Introduction

Skeletal muscle is necessary to accomplish fundamental functions such as the maintenance of the body structure, motility and metabolism by storing and consuming energy. Skeletal muscle development is a multistep process in which myogenic cells are committed to proliferating myogenic precursors that then differentiate into myoblasts and myocytes that fuse to form a multinucleated myotube. Several signals are essential for the regulation of skeletal muscle differentiation involving transcription factors, signalling molecules, transduction pathways and epigenetic modifications. Among these, the histone deacetylases (HDACs) are frequently part of the regulatory elements of muscle genes (Sincennes, Brun, & Rudnicki, n.d.). The HDAC family comprises at least 18 different enzymes classified in four classes in mammals, and has been originally identified for histone deacetylation activity and nucleosome stability. Recent evidence pinpoints their role in deacetylation also of non-histone targets such as p53 and alpha-tubulin (de Leval et al., 2006) as well as in gene transcription (Grunstein, 1997; Megee, Morgan, Mittman, & Smith, 1990). Skeletal muscle is necessary to accomplish fundamental functions such as the maintenance of the body structure, motility and metabolism by storing and consuming energy. Skeletal muscle development is a multistep process in which myogenic cells are committed to proliferating myogenic precursors that then differentiate into myoblasts and myocytes that fuse to form a multinucleated myotube. Several signals are essential for the regulation of skeletal muscle differentiation involving transcription factors, signalling molecules, transduction pathways and epigenetic modifications. Among these, the histone deacetylases (HDACs) are frequently part of the regulatory elements of muscle genes (Sincennes, Brun, & Rudnicki, 2016). The HDAC family comprises at least 18 different enzymes classified in four classes in mammals, and has been originally identified for histone deacetylation activity and nucleosome stability. Recent evidence pinpoints their role in deacetylation also of non-histone targets such as p53 and alpha-tubulin (de Leval et al., 2006) as well as in gene transcription (Grunstein, 1997; Megee et al., 1990).

HDAC8 is the last cloned and characterized member of class I HDACs (Buggy et al., 2000; Van den Wyngaert et al., 2000), it diverges from other class I enzymes as the C-terminal protein-binding domain is not present, probably indicating a functional specialization during evolution (Gregoret, Lee, & Goodson, 2004)(Somoza et al., 2004). HDAC8 is ubiquitously expressed and can localize to either the nucleus or the cytoplasm interacting with non-histone proteins such as the cohesin protein SMC3, estrogen receptor α (ERR α), p53, inv(16) fusion protein (Deardorff et al., 2012; Durst, Lutterbach, Kummalu, Friedman, & Hiebert, 2003; Wilson, Tremblay, Deblois, Sylvain-Drolet, & Giguère, 2010; Wu et al., 2013). Moreover, in normal human tissues *HDAC8* is expressed by smooth muscle including vascular and visceral smooth muscle cells, myoepithelial cells, and myofibroblasts (Durst et al., 2003; Wu et al., 2013) where interacts with cortical actin-binding protein cortactin and Smooth Muscle Actin (SMA) and regulates smooth muscle contraction (Buggy et al., 2000; J. Li et al., 2014; Olson et al., 2014).

In this study, we describe for the first time a specific *HDAC8* expression in human and zebrafish (*Danio rerio*) skeletal muscle and murine and human myogenic cells. In particular, we have analyzed the time course of *HDAC8* expression during skeletal muscle differentiation in murine C2C12 myoblasts and zebrafish. We noticed that *HDAC8* is mainly expressed when differentiation is already started; moreover, in rhabdomyosarcoma derived cell lines RD/12 and RD/18 with low and high differentiation potency respectively, the increment of *HDAC8* expression during the differentiation is prominent in RD/18 than in RD/12 cell line. We also demonstrate that HDAC8 promotes muscle differentiation *in vitro* and *in vivo* as the pharmacological block of its deacetylase activity inhibits myogenesis in the C2C12 cellular model and in zebrafish. This function is accomplished through the canonical Wnt pathway that is down-regulated when HDAC8 activity is inhibited. Our results link for the first time the HDAC8 activity to broad aspects of skeletal muscle development and open new possibility in the use of HDAC8 specific inhibitors (*i.e.* PCI-34051) (Balasubramanian et al., 2008)) for therapeutic intervention on skeletal muscle diseases.

Material and Methods

Animals

Zebrafish (*Danio rerio*) embryos were raised and maintained under standard conditions and national guidelines (Italian decree 4th March 2014, n.26). All experimental procedures were approved by IACUC (Institutional Animal Care and Use Committee). Zebrafish AB strains obtained from the Wilson lab, University College London, London, United Kingdom were maintained at 28°C on a 14 h light/10 h dark cycle. Embryos were collected by natural spawning, staged according to Kimmel and colleagues (Kimmel, Ballard, Kimmel, Ullmann, & Schilling, 1995) and raised at 28°C in fish water (Instant Ocean, 0,1% Methylene Blue in Petri dishes), according to established techniques. We express the embryonic ages in hours post fertilization (hpf) and days post fertilization (dpf). After 24 hpf, to prevent pigmentation 0,003% 1-phenyl-2-thiourea (Sigma-Aldrich, Saint Louis, Missouri, USA) was added to the fish water. Embryos were washed, dechorionated and anaesthetized, with 0.016% tricaine (Ethyl 3-aminobenzoate methanesulfonate salt; Sigma-Aldrich), before observations and picture acquisitions. Embryos were fixed overnight in 4% paraformaldehyde (Sigma-Aldrich) in PBS at 4 °C, then dehydrated stepwise to methanol and stored at -20 °C.

C2C12 and rhabdomyosarcoma cells

C2C12 cells were maintained in growth medium Dulbecco's modified Eagle medium (DMEM) supplemented with 10% fetal bovine serum (FBS, Euroclone, Pero, Italy), 100 IU/mL penicillin and 100 µg/mL streptomycin in a humidified incubator at 37 °C with 5% CO₂. After reaching 80–90% confluence, cells were washed in phosphate-buffered saline (PBS) and differentiated in DMEM medium with Horse Serum 2% (HS, Thermo Fisher Scientific, Waltham, MS, USA). The medium was changed every 48 hours and cultured up to 9 days of differentiation.

RD/12 and RD/18 cell lines were two different clone originally isolated from the human embryonal rhabdomyosarcoma cell lines RD by Lollini and colleagues (Lollini et al., 1991). Cells were cultured in DMEM supplemented with 100 IU/mL penicillin, 100 µg/mL streptomycin and either 10% fetal bovine serum or 2% horse serum. The culture medium was renewed every 48-72 hours up to 11 days of culture in differentiation medium.

RT-PCR and quantitative real time PCR (qPCR)

Total RNAs were isolated from C2C12, RD/12, RD/18 cells and zebrafish embryos at different developmental stages using Trizol reagent (Life Technologies, Carlsbad, CA, USA) according to the producer's instructions. After treatment with DNase I RNase-free (Roche, Basel, Switzerland) to avoid possible genomic contamination, 1µg of RNA was reverse-transcribed using the "ImProm-II™ Reverse Transcription System" (Promega, Madison, WI, USA) and a mixture of oligo(dT) and random primers according to manufacturer's instructions. qPCRs on C2C12 and rhabdomyosarcoma RNAs were carried out in a total volume of 20 µl containing 1X SsoAdv Universal SYBR Green Super Mix (Bio-Rad, Hercules, CA, USA), using proper amount of the RT reaction. qPCRs were performed using the CFX-96™ (Bio-Rad). Relative expression of *HDAC8* was normalized with different reference genes, in particular TATA-box binding protein (TBP) and glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was used for C2C12 cell line while Actin and beta-2-microglobulin (B2M) for rhabdomyosarcoma cell lines.

qPCRs in zebrafish were carried out in a total volume of 20 µl containing 1X iQ SYBR Green Super Mix (Promega), using proper amount of the RT reaction. PCRs were performed using the BioRad iCycler iQ Real Time Detection System (BioRad). For normalization purposes, *rpl8* expression levels were tested in parallel with the gene of interest. Primer list in Supplementary Table 1.

In situ hybridization, histological analysis and immunohistochemistry

Whole mount *in situ* hybridization (WISH) experiments, were carried out as described by Thisse and colleagues (Thisse & Thisse, 2008). Antisense riboprobes were previously *in vitro* labelled with modified nucleotides (*i.e.* digoxigenin, fluorescein, Roche). *hdac8* probe was cloned in our laboratory. Primer list in Supplementary Table 1. WISH experiments were done at least in 3 batches of embryos of (minimum 30 embryos for each category). Immunohistochemistry analysis was carried out on 6 um-thick cryosections from human skeletal muscle biopsy. The muscle biopsy was performed after informed consent, snap-frozen in isopentane/liquid nitrogen, and maintained in liquid nitrogen. Cryosections were permeabilized in cold methanol (MetOH) 50% for 1 minute and MetOH 100% for 1 minute. Cryosections were hydrated with PBS and then blocked for 30 min at room temperature in Normal Goat Serum (NGS) 1X and incubated with primary and secondary antibodies. Primary antibodies were anti-HDAC8 (1:100) (polyclonal clone (H-145): sc-11405, Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) and anti-Lamin B (1:100) (monoclonal clone, Novocastra/YLEM, New Castle-upon-Tyne, UK). Secondary antibody were Alexa 488-conjugated goat anti-mouse IgG or Alexa 546-conjugated goat anti-rabbit IgG, (Invitrogen Life Technologies, Carlsbad, CA, USA) both diluted 1:2000. As control, sections were incubated either with isotype specific IgG or the primary antibody was omitted. Sections were examined either under a Zeiss fluorescence microscope. Immunohistochemistry in zebrafish was carried out as described in Pistocchi and colleagues (Pistocchi, Gaudenzi, et al., 2013). Primary antibody was mouse anti-sarcomeric (MF20, DSHB, diluition 1:4). Secondary antibody was EnVision+ System-HRP Labelled Polymer anti-mouse (Dako, Glostrup, Denmark). Images of embryos and sections were acquired using a microscope equipped with a digital camera with LAS Leica imaging software (Leica, Wetzlar, Germany). Images were processed using the Adobe Photoshop software and when necessary, different focal images planes of the same image have been took separately and later merged in a single image.

Injectons

1
2
3 1 Injections were carried out on 1- to 2-cell stage embryos; the dye tracer rhodamine dextran was also
4
5 2 co-injected. To repress *hdac8* mRNA translations, one morpholino was synthesized (Gene Tools
6
7 3 LLC, Philomath OR, USA) targeting *hdac8*-ATG. and used at the concentration of 1 pmole/embryo
8
9 4 in 1x Danieau buffer (pH 7.6). A standard control morpholino oligonucleotide (ctrl-MO) was
10
11 5 injected in parallel (Nasevicius & Ekker, 2000). ATG-*hdac8*-MO: 5'-
12
13 6 CATTACTGTCGCTTTTTTCACTCAT-3'.

7 8 ***PCI-34051 treatment***

9 For C2C12 cells, HDAC8 inhibitor PCI-34051 (PCI) (Cayman Chemical; Ann Arbor, MI, USA)
10 was administrated at 25 μ M together with differentiating-medium; negative controls were treated
11 with the solvent Dimethyl-sulfoxide (DMSO). The PCI was changed every 24 hours until myogenic
12 differentiation. Zebrafish embryos after the shield developmental stage (6 hpf), were treated with
13 150 μ M PCI added to the fish water at 28°C kept in dark. As a control DMSO was used at the same
14 concentration. The PCI was changed every 24 hours and the embryos are let grown until the desired
15 developmental stage. For dose-dependent assays in zebrafish, the PCI was administrated at 50, 100,
16 150, 250 μ M

17 18 ***Western Blotting***

19 Whole cell extracts from at least 30 zebrafish embryos were classically prepared in RIPA buffer (50
20 mM Tris-HCl pH 7.4, 1% NP-40, 150 mM NaCl, 0.25% sodium deoxycholate, 1mM EDTA, 1mM
21 PMSF, protease inhibitors Roche) (2 μ l/embryo or 1 μ l/ tail). Yolk was previously removed from
22 embryos to avoid yolk protein contamination. The protein concentration was determined using a
23 Micro BCA protein assay kit according to the manufacturer's instructions (Euroclone). 30-40 μ g of
24 each sample were loaded onto a 7.5% or 10% polyacrylamide gels and subjected to electrophoresis.
25 The proteins were then transferred onto PVDF membranes which were blocked using a blocking

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 solution at room temperature for 1 hour prior to incubation with the primary antibodies listed in
2 Supplementary Table 2. After incubation with the HRP-conjugated secondary antibodies for 1 h at
3 room temperature (Secondary antibodies are listed in Supplementary Table 2). The protein bands
4 were detected using ECL detection systems. Imaging acquisition has been done with the Alliance
5 MINI HD9 AUTO Western Blot Imaging System (UVItec Limited, Cambridge) and analysed with
6 the related software (Bellipanni, Murakami, & Weinberg, 2010).

For Peer Review

Results

HDAC8 is expressed in skeletal muscle and its expression correlates with an advance differentiated state of muscle cells.

Several expression profiles of *HDAC8* suggested that it has a ubiquitous expression in human tissues, with higher expression in particular organs such as brain, pancreas, kidney, prostate, liver and smooth muscles. *HDAC8* transcript and protein have been detected both in the nucleus and cytosol, suggesting that *HDAC8* might have a variable localization within the cell, depending on the cell type and its post-translational modifications such as phosphorylation (Buggy et al., 2000; de Ruijter, van Gennip, Caron, Kemp, & van Kuilenburg, 2003; Hu et al., 2000; Waltregny et al., 2004). Using immunofluorescence assays we detected for the first time an expression of *HDAC8* in normal human skeletal muscle with a predominant nuclear localization of the protein, as shown by the co-localization of *HDAC8* and Lamin B (Figure 1A-C).

In parallel, we cloned the zebrafish orthologue of human *HDAC8* (Chr 7: 51,656,099-51,710,015), and by whole mount *in situ* hybridization analyses (WISH) we confirmed the expression of *hdac8* in skeletal muscle of zebrafish embryos at different developmental stages (Figure 1D-F'). In zebrafish the expression of *hdac8* varied among the developmental stages analysed (24, 36 and 48 hours post fertilization, hpf), and was increased at 36 hpf when the first myogenic wave have already occurred (Stellabotte, Dobbs-McAuliffe, Fernandez, Feng, & Devoto, 2007) (Figure 1D-F').

We therefore investigated a possible correlation between *HDAC8* expression and skeletal muscle differentiation progression. We first examined its expression in murine C2C12 skeletal myogenic cells, which represent a highly suitable model for analysis of myogenic differentiation. C2C12 myoblasts proliferate in growth medium with high serum concentration (10% FBS) until they reach confluence, while differentiation into multinucleated myotubes is triggered shifting to differentiation medium with low serum concentration (2% horse serum). *Hdac8* transcript, analysed by qRT-PCR techniques, was present in C2C12 cells in growth medium and in differentiation medium at 1 days but its expression was significantly increased at 7 and 9 days of differentiation

(Figure 1G). In zebrafish the expression of *hdac8* analysed by qRT-PCR techniques confirmed the results previously shown by WISH, as the transcript is increased after the first myogenic wave when differentiation was accomplished with an expression peak at 36 hpf (Figure 1H). To further confirm the correlation between *HDAC8* expression and an advanced stage of differentiation, we choose two different subclones of the rhabdomyosarcoma cell line RD which differ in the differentiation potency: the RD/18 cells are able to reach a terminal differentiation while the RD/12 cells do not fully differentiate (Lollini et al., 1991). The expression of *HDAC8* was significantly increased at 11 days of differentiation with an increment of 4 fold in RD/18 and about 1 fold in RD/12 (Figure 1I).

HDAC8 activity regulates skeletal muscle differentiation in zebrafish and C2C12 myoblasts.

To investigate a possible function of HDAC8 in differentiating skeletal muscles we took advantage of the well characterized PCI inhibitor that blocks HDAC8 deacetylase activity (Balasubramanian et al., 2008). We administrated PCI to zebrafish embryos *in-vivo* and C2C12 cells *in-vitro*. Zebrafish embryos were treated with a concentration of 150 μ M of PCI from the 50% stage of epiboly, a developmental stage in which the mesodermal layer, from which skeletal muscle derives, is positioning in the gastrula. At 48 hpf, zebrafish embryos presented morphological defects in the Central Nervous System and muscles, the regions where *hdac8* transcript was more expressed as shown in Figure 1. The PCI treated embryos could be divided in three phenotypical classes based on the severity of the CNS and muscle phenotype: class I showed a phenotype comparable to the control embryos treated with the solvent DMSO, class II presented a mild phenotype and class III presented a severe phenotype (Figure 2A-D, class quantification in E). We performed a dose-response assay demonstrating that the observed phenotypes were correlated to the doses of PCI treatment (Suppl. Figure S1). The sarcomeric myosins, that are expressed in differentiated and functional muscle, were diminished in PCI-treated embryos in comparison to controls analysed by immunohistochemistry and Western blot techniques (Figure 2 F-I). Interestingly, same morphological defects and myosin reduction were obtained in zebrafish embryos injected with the

hdac8 morpholino (*hdac8*-MO) that blocks Hdac8 protein production. These data indicate that the skeletal muscle differentiation impairment was specific due to Hdac8 loss-of-function (Suppl. Figure S2). Moreover, at 24 hpf the embryos treated with PCI did not present myogenic impairment confirming that Hdac8 activity is not necessary during early skeletal muscle differentiation (Suppl. Figure S3).

Also *in-vitro*, PCI treatment blocked differentiation of C2C12 myoblasts in comparison to DMSO treated cells. Under differentiating conditions, wild-type C2C12 cells fused into multinucleated myotubes. By contrast, when challenged to differentiate in low-serum medium in presence of PCI, C2C12 cells remained mononucleated and maintained an undifferentiated phenotype. We assessed that the differentiation of PCI treated cells was impaired in comparison to DMSO treated cells as the levels of sarcomeric myosins analysed by Western blot technique were diminished (Figure 2J-K).

HDAC8 regulates skeletal muscle differentiation through the activation of the canonical Wnt pathway.

In order to gain mechanistic insights into how HDAC8 regulates skeletal muscle differentiation, we hypothesized that it modulates the canonical Wnt pathway, a well-known regulator of skeletal muscle development and differentiation (Rudnicki & Williams, 2015). Indeed, in a hepatocellular model, it has been demonstrated that HDAC8 positively regulates the β -catenin/TCF signalling acting in concert with EZH2 to epigenetically repress Wnt antagonists (Tian et al., 2015). Therefore, we analysed the activation status of the canonical Wnt pathway in zebrafish embryos and C2C12 myoblasts treated with PCI. The phosphorylated and active form of β -catenin was diminished by Western blot analyses in PCI treated zebrafish embryos in comparison to controls treated with the DMSO. By contrast, the levels of total β -catenin were even increased (Figure 3A, quantification in B and C). This last result is not surprising since we have seen similar up-regulation

1
2
3 1 of β -Catenin in zebrafish embryos with impaired activity of the canonical Wnt pathway (Valenti et
4
5 2 al., 2015). To verify the efficiency of the PCI-mediated Hdac8 inhibition that is responsible of the
6
7 3 Wnt pathway down-regulation, we analysed the acetylation status of Smc3, a known Hdac8 target
8
9 4 (Deardorff et al., 2012). Acetylated Smc3 (Smc3ac) levels were increased following PCI treatment
10
11 5 of the embryos, confirming the block of Hdac8 activity (Figure 3D, quantification in E). Same
12
13 6 results were obtained in the C2C12 cells in differentiation medium treated with PCI: Western blot
14
15 7 analyses confirmed the lower expression of active β -catenin in comparison to total β -catenin
16
17 8 (Figure Figure 3F, quantification in G and H) and increased levels of Smc3ac following PCI
18
19 9 treatment (Figure 3I, quantification in J).
20
21
22 10 The Wnt pathway in zebrafish can be activated through chemical treatments such as LiCl
23
24 11 (Pistocchi, Fazio, et al., 2013). Therefore, to further demonstrate that skeletal muscle differentiation
25
26 12 impairment observed with PCI-mediated Hdac8 inhibition was specifically due to Wnt pathway
27
28 13 down-regulation, we re-activated the pathway adding LiCl in PCI-treated zebrafish embryos. The
29
30 14 morphological defects presented by PCI-treated embryos at 36 hpf (embryos with morphological
31
32 15 defects: 50/70) were partially rescued by LiCl addition (embryos with morphological defects:
33
34 16 20/70) (Figure 4A-C). Moreover, the levels of sarcomeric myosins analysed by Western blot
35
36 17 techniques were rescued in embryos treated with PCI+LiCl in comparison to embryos treated only
37
38 18 with PCI (Figure 4D, quantification in E). We also verify the efficiency of LiCl treatment by
39
40 19 measuring the active β -catenin levels (Figure 4D, quantification in F).
41
42
43
44 20
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Discussion

In previous works HDAC8 was shown to be expressed in smooth muscle cells in association with SMA and cortactin (Jia Li et al., 2014) and its silencing by RNA interference (RNAi) impairs the contraction of smooth muscle cultured cells (Waltregny et al., 2005). However, the role and mechanism of HDAC8 action in smooth muscle tissues are largely unknown. In this work, we described for the first time the expression and role of HDAC8 in the skeletal muscle. Firstly, we demonstrated that HDAC8 is expressed in human and zebrafish skeletal muscle; then we analysed the expression of *HDAC8* during muscle differentiation in the murine C2C12 skeletal muscle cells, during zebrafish muscle development and in two types of rhabdomyosarcoma cells with various degree of invasiveness correlating to their ability to differentiate (RD/12 and RD/18). We decided to include these cells in the expression analyses as it has been reported that HDACi synergize with current anticancer drugs to induce apoptosis in rhabdomyosarcoma although the authors observed a switch to myogenic differentiation (Vleeshouwer-Neumann et al., 2015; Di Pompo et al., 2015). Interestingly, we correlate the expression of *HDAC8* with an advanced differentiation state of skeletal muscles. Indeed, both in C2C12 cells and zebrafish, *HDAC8* expression is weak in the initial phases and increases later during the muscle differentiation process. These data are even more striking in the rhabdomyosarcoma cells, where the RD/18 cell line cultured in the differentiation medium for 11 days shows a greater increase in the HDAC8 expression compared to the RD/12 cell line maintained in the same conditions. This increase correlates with the differentiation capacity of the two cell lines.

For functional analyses, we treated the C2C12 cells and the zebrafish embryos with the HDAC8 inhibitor PCI-34051. In zebrafish, we also performed loss-of-function studies by injecting the oligonucleotide antisense morpholino designed against *hdac8* to compare and confirm the results obtained with the PCI-34051 treatment. Both in the cellular and zebrafish models with reduced HDAC8 activity, we observed an impairment in muscle differentiation following the initial myoblast commitment, in line with the kinetic of *HDAC8* expression previously analysed. In the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

C2C12 cells, myoblasts were formed but failed to fuse in myotubes and to express the markers of differentiation; in zebrafish, the levels of functional myosins were reduced after 24 hpf but the myogenic program started, as demonstrated by the proper expression of the MRFs MyoD and Myog and by the presence of myosin proteins. Interestingly, it has been already shown that the levels of myogenin were not affected by myoblast exposure to HDACi (Iezzi, Cossu, Nervi, Sartorelli, & Puri, 2002), suggesting that HDACi selectively activate late muscle markers. It is also reported a dual action for HDACi on muscle differentiation, depending on the stage of administration: previous studies reported that HDACi have different effects by promoting or inhibiting myogenesis (Steinbach, Wolffe, & Rupp, 1997) and this discrepancy might be explained by the stage-specific effects of HDACi exposure. In zebrafish embryos, we performed the Hdac8 inhibition by adding the PCI-34051 inhibitor after the shield stage (6 hpf) to prevent gross morphological defects in the initial phase of gastrulation when mesoderm is defined.

The block on muscle differentiation observed following HDAC8 inhibition is correlated with the down-regulation of the canonical Wnt pathway. Several works demonstrate that the formation of skeletal muscle is tightly modulated by Wnt signalling for self-renewal and muscle differentiation and its dysregulation leads to perturbation of muscle fibers. Chemical modulation of the Wnt/ β -catenin pathway in differentiating myoblasts, using the activator LiCl, increases both the number and size of C2C12 myotubes while inhibitors of Wnt/ β -catenin signalling result in a significant decrease in myotube length (Abraham, 2016). Indeed, the Wnt target β -catenin interacts directly with MyoD, enhancing its binding to E box elements and its transcriptional activity of muscle specific genes. This transactivation is inhibited when β -catenin is deficient or the interaction between MyoD and β -catenin is disrupted (Kim, Mei 2008). We demonstrate that the reduction of myosins observed in PCI-34051 treated embryos was caused by a decrease in activated β -catenin levels. A mechanism by which HDAC8 regulates the canonical Wnt pathway has been recently described in human NAFLD-associated hepatocellular carcinoma (HCC) by Tian and colleagues

(Tian, Mok, Yang, & Cheng, 2016). HDAC8 physically interacts with the polycomb protein enhancer of zeste homolog 2 (EZH2) and contributes to the activation of Wnt/ β -catenin signalling. Further analyses are necessary to demonstrate whether this mechanism is conserved also in skeletal muscle and acetylome profile following PCI inhibition may uncover HDAC8-related targets. Dis-regulation of canonical Wnt signalling has been reported in different muscle pathologies, such as Duchenne Muscular Dystrophy (DMD) (Trensz, Haroun, Cloutier, Richter, & Grenier, 2010), FascioScapuloHumeral Muscular Dystrophy (FSHD) (Block et al., 2013) and OculoPharyngeal Muscular Dystrophy (OPMD) (Abu-Baker et al., 2013). Inhibition of canonical Wnt signalling by Dkk in a mouse model for DMD (mdx), was shown to reduce fibrosis (Trensz et al., 2010). HDACi are recently emerged as potential pharmacological strategies for cancer treatment, and several of them are already approved by the international Drug Administration agencies. The increasing interest and use of HDACi has led to the development of class-specific inhibitors, such as the PCI-34051, which helps us to uncover the functional role of HDAC8 in skeletal muscle differentiation and, in the future, might ameliorate the phenotype in pathological conditions. Based on the numerous beneficial effects of HDACi in skeletal muscle under pathological conditions, we believe that they are promising therapeutics.

Competing interests

All authors declare that they have no conflict of interest. Declaration of interest: none.

Author contributions: conceived and designed the experiments: AP¹, AM¹, PV¹. Performed the experiments on human samples: CB^{2,5}, MM². Performed the experiments in zebrafish: MS¹, SE¹, LB¹, AP¹, AP¹, LF¹. Performed the experiments in C2C12 cells: LF¹, AMF^{4,6}, FP³, FF³. Performed the experiments in rhabdomyosarcoma cells: FP³, FF³. Analyzed the data on human sample: CB^{2,5}, MM². Analysed the data in zebrafish: MS¹, SE¹, LB¹, AP¹, AP¹, LF¹, PR¹, AG^{7,8}, GB^{7,8}. Analysed the data in C2C12 cells: LF¹, PR¹, FP³, FF³, FB, AMF^{4,6}, GC⁴. Wrote the paper: AP¹. Supervised paper drafting: AP¹, AG^{7,8}, GB^{7,8}. Supervised the research project AP¹.

References

- Abraham, S. T. (2016). A role for the Wnt3a/ β -catenin signaling pathway in the myogenic program of C2C12 cells. *In Vitro Cellular and Developmental Biology - Animal*, 52(9), 935–941. <https://doi.org/10.1007/s11626-016-0058-5>
- Abu-Baker, A., Laganiere, J., Gaudet, R., Rochefort, D., Brais, B., Neri, C., ... Rouleau, G. A. (2013). Lithium chloride attenuates cell death in oculopharyngeal muscular dystrophy by perturbing Wnt/ β -catenin pathway. *Cell Death & Disease*, 4(10), e821. <https://doi.org/10.1038/cddis.2013.342>
- Balasubramanian, S., Ramos, J., Luo, W., Sirisawad, M., Verner, E., & Buggy, J. J. (2008). A novel histone deacetylase 8 (HDAC8)-specific inhibitor PCI-34051 induces apoptosis in T-cell lymphomas. *Leukemia*, 22(5), 1026–1034. <https://doi.org/10.1038/leu.2008.9>
- Bellipanni, G., Murakami, T., & Weinberg, E. S. (2010). Molecular dissection of Otx1 functional domains in the zebrafish embryo. *Journal of Cellular Physiology*, 222(2), 286–293. <https://doi.org/10.1002/jcp.21944>
- Block, G. J., Narayanan, D., Amell, A. M., Petek, L. M., Davidson, K. C., Bird, T. D., ... Miller, D. G. (2013). Wnt/ β -catenin signaling suppresses DUX4 expression and prevents apoptosis of FSHD muscle cells. *Human Molecular Genetics*, 22(23), 4661–72. <https://doi.org/10.1093/hmg/ddt314>
- Buggy, J. J., Sideris, M. L., Mak, P., Lorimer, D. D., McIntosh, B., & Clark, J. M. (2000). Cloning and characterization of a novel human histone deacetylase, HDAC8. *The Biochemical Journal*, 350 Pt 1, 199–205. <https://doi.org/10.1042/BJ3500199>
- de Leval, L., Waltregny, D., Boniver, J., Young, R. H., Castronovo, V., & Oliva, E. (2006). Use of histone deacetylase 8 (HDAC8), a new marker of smooth muscle differentiation, in the classification of mesenchymal tumors of the uterus. *The American Journal of Surgical Pathology*, 30(3), 319–27. <https://doi.org/10.1097/01.pas.0000188029.63706.31>
- de Ruijter, A. J. M., van Gennip, A. H., Caron, H. N., Kemp, S., & van Kuilenburg, A. B. P. (2003).

1
2
3 1 Histone deacetylases (HDACs): characterization of the classical HDAC family. *The*
4
5 2 *Biochemical Journal*, 370(Pt 3), 737–49. <https://doi.org/10.1042/BJ20021321>
6
7 3 Deardorff, M. A., Bando, M., Nakato, R., Watrin, E., Itoh, T., Minamino, M., ... Shirahige, K.
8
9 4 (2012). HDAC8 mutations in Cornelia de Lange syndrome affect the cohesin acetylation cycle.
10
11 5 *Nature*, 489(7415), 313–317. <https://doi.org/10.1038/nature11316>
12
13 6 Di Pompo, G., Salerno, M., Rotili, D., Valente, S., Zwergel, C., Avnet, S., ... Mai, A. (2015). Novel
14
15 7 Histone Deacetylase Inhibitors Induce Growth Arrest, Apoptosis, and Differentiation in
16
17 8 Sarcoma Cancer Stem Cells. *Journal of Medicinal Chemistry*, 58(9), 4073–4079.
18
19 9 <https://doi.org/10.1021/acs.jmedchem.5b00126>
20
21
22 10 Durst, K. L., Lutterbach, B., Kummalue, T., Friedman, A. D., & Hiebert, S. W. (2003). The inv(16)
23
24 11 fusion protein associates with corepressors via a smooth muscle myosin heavy-chain domain.
25
26 12 *Molecular and Cellular Biology*, 23(2), 607–19. [https://doi.org/10.1128/MCB.23.2.607–](https://doi.org/10.1128/MCB.23.2.607-619.2003)
27
28 13 619.2003
29
30
31 14 Gregoret, I. V., Lee, Y. M., & Goodson, H. V. (2004). Molecular evolution of the histone
32
33 15 deacetylase family: Functional implications of phylogenetic analysis. *Journal of Molecular*
34
35 16 *Biology*, 338(1), 17–31. <https://doi.org/10.1016/j.jmb.2004.02.006>
36
37 17 Grunstein, M. (1997). Histone acetylation in chromatin structure and transcription. *Nature*,
38
39 18 389(6649), 349–352. <https://doi.org/10.1038/38664>
40
41
42 19 Hu, E., Chen, Z., Fredrickson, T., Zhu, Y., Kirkpatrick, R., Zhang, G. F., ... Winkler, J. (2000).
43
44 20 Cloning and characterization of a novel human class I histone deacetylase that functions as a
45
46 21 transcription repressor. *Journal of Biological Chemistry*, 275(20), 15254–15264.
47
48 22 <https://doi.org/10.1074/jbc.M908988199>
49
50 23 Iezzi, S., Cossu, G., Nervi, C., Sartorelli, V., & Puri, P. L. (2002). Stage-specific modulation of
51
52 24 skeletal myogenesis by inhibitors of nuclear deacetylases. *Proceedings of the National*
53
54 25 *Academy of Sciences of the United States of America*, 99(11), 7757–62.
55
56 26 <https://doi.org/10.1073/pnas.112218599>
57
58
59
60

- 1 Kimmel, C., Ballard, W., Kimmel, S., Ullmann, B., & Schilling, T. (1995). Stages of embryonic
2 development of the zebrafish. *Developmental Dynamics*, 203(3), 253–310.
3 <https://doi.org/10.1002/aja.1002030302>
- 4 Li, J., Chen, S., Cleary, R. A., Wang, R., Gannon, O. J., Seto, E., & Tang, D. D. (2014). Histone
5 deacetylase 8 regulates cortactin deacetylation and contraction in smooth muscle tissues.
6 *American Journal of Physiology. Cell Physiology*, 307(3), C288–95.
7 <https://doi.org/10.1152/ajpcell.00102.2014>
- 8 Li, J., Chen, S., Cleary, R. A., Wang, R., Gannon, O. J., Seto, E., & Tang, D. D. (2014). Histone
9 deacetylase 8 regulates cortactin deacetylation and contraction in smooth muscle tissues. *AJP:*
10 *Cell Physiology*, 307(3), C288–C295. <https://doi.org/10.1152/ajpcell.00102.2014>
- 11 Lollini, P. L., De Giovanni, C., Landuzzi, L., Nicoletti, G., Scotlandi, K., & Nanni, P. (1991).
12 Reduced metastatic ability of in vitro differentiated human rhabdomyosarcoma cells. *Invasion*
13 *& Metastasis*, 11(2), 116–24. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/1917385>
- 14 Megee, P. C., Morgan, B. A., Mittman, B. A., & Smith, M. M. (1990). Genetic analysis of histone
15 H4: Essential role of lysines subject to reversible acetylation. *Science*, 247(4944), 841–845.
16 <https://doi.org/10.1126/science.2106160>
- 17 Nasevicius, a, & Ekker, S. C. (2000). Effective targeted gene “knockdown” in zebrafish. *Nature*
18 *Genetics*, 26(2), 216–20. <https://doi.org/10.1038/79951>
- 19 Olson, D. E., Udeshi, N. D., Wolfson, N. A., Pitcairn, C. A., Sullivan, E. D., Jaffe, J. D., ... Holson,
20 E. B. (2014). An unbiased approach to identify endogenous substrates of “histone” deacetylase
21 8. *ACS Chemical Biology*, 9(10), 2210–2216. <https://doi.org/10.1021/cb500492r>
- 22 Pistocchi, A., Fazio, G., Cereda, A., Ferrari, L., Bettini, L. R., Messina, G., ... Massa, V. (2013).
23 Cornelia de Lange Syndrome: NIPBL haploinsufficiency downregulates canonical Wnt
24 pathway in zebrafish embryos and patients fibroblasts. *Cell Death and Disease*, 4(10), e866.
25 <https://doi.org/10.1038/cddis.2013.371>
- 26 Pistocchi, A., Gaudenzi, G., Foglia, E., Monteverde, S., Moreno-Fortuny, A., Pianca, A., ...

1
2
3 1 Messina, G. (2013). Conserved and divergent functions of Nfix in skeletal muscle
4
5 2 development during vertebrate evolution. *Development (Cambridge)*, 140(7).
6
7 3 <https://doi.org/10.1242/dev.076315>
8
9 4 Rudnicki, M. A., & Williams, B. O. (2015). Wnt signaling in bone and muscle. *Bone*.
10
11 5 <https://doi.org/10.1016/j.bone.2015.02.009>
12
13 6 Sincennes, M.-C., Brun, C. E., & Rudnicki, M. A. (n.d.). Tissue-Specific Progenitor and Stem Cells
14
15 7 Concise Review: Epigenetic Regulation of Myogenesis in Health and Disease.
16
17 8 <https://doi.org/10.5966/sctm.2015-0266>
18
19 9 Sincennes, M.-C., Brun, C. E., & Rudnicki, M. A. (2016). Concise Review: Epigenetic Regulation
20
21 10 of Myogenesis in Health and Disease. *STEM CELLS Translational Medicine*, 5(3), 282–290.
22
23 11 <https://doi.org/10.5966/sctm.2015-0266>
24
25 12 Somoza, J. R., Skene, R. J., Katz, B. A., Mol, C., Ho, J. D., Jennings, A. J., ... Tari, L. W. (2004).
26
27 13 Structural snapshots of human HDAC8 provide insights into the class I histone deacetylases.
28
29 14 *Structure*, 12(7), 1325–1334. <https://doi.org/10.1016/j.str.2004.04.012>
30
31 15 Steinbach, O. C., Wolffe, A. P., & Rupp, R. A. W. (1997). Somatic linker histones cause loss of
32
33 16 mesodermal competence in *Xenopus*. *Nature*, 389(6649), 395–399.
34
35 17 <https://doi.org/10.1038/38755>
36
37 18 Stellabotte, F., Dobbs-McAuliffe, B., Fernandez, D. A., Feng, X., & Devoto, S. H. (2007). Dynamic
38
39 19 somite cell rearrangements lead to distinct waves of myotome growth. *Development*, 134(7),
40
41 20 1253–1257. <https://doi.org/10.1242/dev.000067>
42
43 21 Thisse, C., & Thisse, B. (2008). High-resolution in situ hybridization to whole-mount zebrafish
44
45 22 embryos. *Nature Protocols*, 3(1), 59–69. <https://doi.org/10.1038/nprot.2007.514>
46
47 23 Tian, Y., Mok, M., Yang, P., & Cheng, A. (2016). Epigenetic Activation of Wnt/ β -Catenin
48
49 24 Signaling in NAFLD-Associated Hepatocarcinogenesis. *Cancers*, 8(8), 76.
50
51 25 <https://doi.org/10.3390/cancers8080076>
52
53 26 Tian, Y., Wong, V. W. S., Wong, G. L. H., Yang, W., Sun, H., Shen, J., ... Chan, H. L. Y. (2015).
54
55
56
57
58
59
60

- 1
2
3 1 Histone deacetylase HDAC8 promotes insulin resistance and β -catenin activation in NAFLD-
4 associated hepatocellular carcinoma. *Cancer Research*, 75(22), 4803–4816.
5 2
6
7 3 <https://doi.org/10.1158/0008-5472.CAN-14-3786>
8
9 4 Trenszt, F., Haroun, S., Cloutier, A., Richter, M. V., & Grenier, G. (2010). A muscle resident cell
10 population promotes fibrosis in hindlimb skeletal muscles of mdx mice through the Wnt
11 canonical pathway. *American Journal of Physiology-Cell Physiology*, 299(5), C939–C947.
12 6
13
14 7 <https://doi.org/10.1152/ajpcell.00253.2010>
15
16
17 8 Valenti, F., Ibetti, J., Komiya, Y., Baxter, M., Lucchese, A. M., Derstine, L., ... Bellipanni, G.
19 (2015). The Increase in Maternal Expression of *axin1* and *axin2* Contribute to the Zebrafish
20 Mutant *Ichabod* Ventralized Phenotype. *Journal of Cellular Biochemistry*, 116(3), 418–430.
21
22 10 <https://doi.org/10.1002/jcb.24993>
23
24
25 11
26 12 Van den Wyngaert, I., de Vries, W., Kremer, a, Neefs, J., Verhasselt, P., Luyten, W. H., & Kass, S.
27 U. (2000). Cloning and characterization of human histone deacetylase 8. *FEBS Letters*, 478(1–
28 2), 77–83. [https://doi.org/10.1016/S0014-5793\(00\)01813-5](https://doi.org/10.1016/S0014-5793(00)01813-5)
29 14
30
31 15 Vleeshouwer-Neumann, T., Phelps, M., Bammler, T. K., MacDonald, J. W., Jenkins, I., & Chen, E.
32 Y. (2015). Histone Deacetylase Inhibitors Antagonize Distinct Pathways to Suppress
33 Tumorigenesis of Embryonal Rhabdomyosarcoma. *PloS One*, 10(12), e0144320.
34
35 16
36 17 <https://doi.org/10.1371/journal.pone.0144320>
37
38
39 18
40
41 19 Waltregny, D., De Leval, L., Glénisson, W., Ly Tran, S., North, B. J., Bellahcène, A., ...
42
43 Castronovo, V. (2004). Expression of histone deacetylase 8, a class I histone deacetylase, is
44 restricted to cells showing smooth muscle differentiation in normal human tissues. *The*
45
46 21 *American Journal of Pathology*, 165(2), 553–564. [https://doi.org/10.1016/S0002-](https://doi.org/10.1016/S0002-9440(10)63320-2)
47
48 22
49 23
50 9440(10)63320-2
51
52 24 Waltregny, D., Glénisson, W., Tran, S. L., North, B. J., Verdin, E., Colige, A., & Castronovo, V.
53
54 (2005). Histone deacetylase HDAC8 associates with smooth muscle alpha-actin and is
55
56 25 essential for smooth muscle cell contractility. *Faseb*, 19(8), 966–8.
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 <https://doi.org/10.1096/fj.04-2303fje>
2 Wilson, B. J., Tremblay, A. M., Deblois, G., Sylvain-Drolet, G., & Giguère, V. (2010). An
3 acetylation switch modulates the transcriptional activity of estrogen-related receptor alpha.
4 *Molecular Endocrinology*, 24(7), 1349–1358. <https://doi.org/10.1210/me.2009-0441>
5 Wu, J., Du, C., Lv, Z., Ding, C., Cheng, J., Xie, H., ... Zheng, S. (2013). The up-regulation of
6 histone deacetylase 8 promotes proliferation and inhibits apoptosis in hepatocellular
7 carcinoma. *Digestive Diseases and Sciences*, 58(12). [https://doi.org/10.1007/s10620-013-](https://doi.org/10.1007/s10620-013-2867-7)
8 2867-7

For Peer Review

Figures legends

Figure 1: HDAC8 is expressed in human, murine and zebrafish skeletal muscle and its expression correlates with differentiation potency. (A-C) HDAC8 protein expression in normal human skeletal muscles. Immunofluorescence staining of HDAC8 (A), Lamin B (B) and merge of the two signals (C). The localization of HDAC8 in human skeletal muscle is predominantly nuclear as shown by the co-localization with the Lamin B protein. (D-F) *hdac8* mRNA expression in zebrafish. WISH analyses of *hdac8* transcript localization in skeletal muscle of zebrafish embryos at 24 hpf (D), 36 hpf (E) and 48 hpf (F). Transverse histological sections of the previously hybridized embryos show the localization of *hdac8* transcript in the myotome (D',E',F'). (G) *Hdac8* qRT-PCR analyses on murine C2C12 myoblasts at different stages of differentiation. *Hdac8* expression is increased at 7-9 days after the induction of the differentiation when differentiation is accomplished. (H) *hdac8* qRT-PCR analyses on RNA from 24, 36 and 48 hpf zebrafish embryos. *hdac8* expression is increased at 36 hpf when the first myogenic wave is completed. (I) *HDAC8* qRT-PCR analyses on RD/12 and RD/18 rhabdomyosarcoma cells. At 11 days after the induction of differentiation, *HDAC8* is more expressed in RD/18 cells that are able to fully differentiate in comparison to RD/12 cells. Scale bar represents 50 μ m in (A-C) and 100 μ m (D-F'). Asterisks represent ** $p < 0.01$, *** $p < 0.001$, Student's t test.

Figure 2: Inhibition of HDAC8 activity reduces skeletal muscle differentiation in zebrafish embryos and murine C2C12 myoblasts. (A-E) *In-vivo* treatment of zebrafish embryos with DMSO or PCI. Different phenotypical classes with increased severity (B-D; quantification in E) with PCI treatment compared to the control embryos treated with the DMSO (A). (F-G) Immunohistochemical staining (IHC) and (H-I) western blot analyses of sarcomeric myosins. Sarcomeric myosins are reduced in PCI treated embryos at 48 hpf (G) in comparison to controls (F). Western blot analyses (H; quantification in I) confirmed all MyHC reduction in PCI treated embryos in comparison to controls. (J-K). Inhibition of HDAC8 activity impaired C2C12

differentiation. Western blot analyses (*J*; quantification in *K*) confirmed all MyHC reduction in PCI treated C2C12 cells in comparison to DMSO treated. Scale bars indicates 100 μ m in (*A*, *F*). Asterisks represent **p*<0.05, Student's *t* test.

Figure 3: HDAC8 activates canonical Wnt signalling. (*A-C*) Canonical Wnt signalling was decreased with the PCI treatment in zebrafish embryos. (*A*) Active β -catenin was decreased in PCI treated embryos in comparison to DMSO controls while total β -catenin was increased by Western blot analyses and relative quantifications (*B-C*). (*D-E*) The efficacy of PCI treatment was verified by the acetylation status of the Hdac8 target Smc3. (*D*) Smc3ac levels were increased in PCI treated embryos in comparison to DMSO controls, quantification in (*E*). (*F-H*) Canonical Wnt signalling was decreased with the PCI treatment in C2C12 cells in differentiation medium. (*F*) Active β -catenin was decreased in PCI treated C2C12 cells in comparison to those treated with DMSO, while total β -catenin was increased by Western blot analyses and relative quantifications (*G-H*). (*I-J*) The efficacy of PCI treatment in the C2C12 was verify by the acetylation status of the Hdac8 target Smc3. (*I*) Smc3ac levels were increased in PCI treated C2C12 in comparison to DMSO controls, quantification in (*J*). Asterisks represent **p*<0.05, ****p*<0.001, Student's *t* test.

Figure 4: The HDAC8-mediated positive regulation of Wnt signalling is responsible for skeletal muscle differentiation. (*A-C*) Morphological defect presented by PCI-treated embryos were rescued by LiCl addition. (*D-F*) Skeletal muscle differentiation was rescued when the Wnt pathway was restored by LiCl in PCI treated zebrafish embryos. (*D*) Sarcomeric myosins, analysed by Western blot techniques, decreased in PCI treated embryos and returned comparable to those treated with DMSO when Wnt pathway was rescued adding LiCl (quantification in *E*). The efficacy of LiCl treatment was verify measuring the active β -catenin by Western blot techniques (quantification in *F*). Asterisks represent **p*<0.05, Student's *t* test.

1

2

For Peer Review




Figure 1: HDAC8 is expressed in human, murine and zebrafish skeletal muscle and its expression correlates with differentiation potency. (A-C) HDAC8 protein expression in normal human skeletal muscles. Immunofluorescence staining of HDAC8 (A), Lamin B (B) and merge of the two signals (C). The localization of HDAC8 in human skeletal muscle is predominantly nuclear as shown by the co-localization with the Lamin B protein. (D-F) *hdac8* mRNA expression in zebrafish. WISH analyses of *hdac8* transcript localization in skeletal muscle of zebrafish embryos at 24 hpf (D), 36 hpf (E) and 48 hpf (F). Transverse histological sections of the previously hybridized embryos show the localization of *hdac8* transcript in the myotome (D',E',F'). (G) *Hdac8* qRT-PCR analyses on murine C2C12 myoblasts at different stages of differentiation. *Hdac8* expression is increased at 7-9 days after the induction of the differentiation when differentiation is accomplished. (H) *hdac8* qRT-PCR analyses on RNA from 24, 36 and 48 hpf zebrafish embryos. *hdac8* expression is increased at 36 hpf when the first myogenic wave is completed. (I) HDAC8 qRT-PCR analyses on RD/12 and RD/18 rhabdomyosarcoma cells. At 11 days after the induction of differentiation, HDAC8 is more expressed in RD/18 cells that are able to fully differentiate in comparison to RD/12 cells. Scale bar represents 50 μ m in (A-C) and 100 μ m (D-F'). Asterisks represent **p<0.01, ***p<0.001, Student's t test.

216x202mm (300 x 300 DPI)




Figure 2: Inhibition of HDAC8 activity reduces skeletal muscle differentiation in zebrafish embryos and murine C2C12 myoblasts. (A-E) In-vivo treatment of zebrafish embryos with DMSO or PCI. Different phenotypical classes with increased severity (B-D; quantification in E) with PCI treatment compared to the control embryos treated with the DMSO (A). (F-G) Immunohistochemical staining (IHC) and (H-I) western blot analyses of sarcomeric myosins. Sarcomeric myosins are reduced in PCI treated embryos at 48 hpf (G) in comparison to controls (F). Western blot analyses (H; quantification in I) confirmed all MyHC reduction in PCI treated embryos in comparison to controls. (J-K). Inhibition of HDAC8 activity impaired C2C12 differentiation. Western blot analyses (J; quantification in K) confirmed all MyHC reduction in PCI treated C2C12 cells in comparison to DMSO treated. Scale bars indicates 100 μ m in (A, F). Asterisks represent * $p < 0.05$, Student's t test.

588x233mm (96 x 96 DPI)




Figure 3: HDAC8 activates canonical Wnt signalling. (A-C) Canonical Wnt signalling was decreased with the PCI treatment in zebrafish embryos. (A) Active β -catenin was decreased in PCI treated embryos in comparison to DMSO controls while total β -catenin was increased by Western blot analyses and relative quantifications (B-C). (D-E) The efficacy of PCI treatment was verified by the acetylation status of the Hdac8 target Smc3. (D) Smc3ac levels were increased in PCI treated embryos in comparison to DMSO controls, quantification in (E). (F-H) Canonical Wnt signalling was decreased with the PCI treatment in C2C12 cells in differentiation medium. (F) Active β -catenin was decreased in PCI treated C2C12 cells in comparison to those treated with DMSO, while total β -catenin was increased by Western blot analyses and relative quantifications (G-H). (I-J) The efficacy of PCI treatment in the C2C12 was verify by the acetylation status of the Hdac8 target Smc3. (I) Smc3ac levels were increased in PCI treated C2C12 in comparison to DMSO controls, quantification in (J). Asterisks represent *p<0.05, ***p<0.001, Student's t test.




Figure 4: The HDAC8-mediated positive regulation of Wnt signalling is responsible for skeletal muscle differentiation. (A-C) Morphological defect presented by PCI-treated embryos were rescued by LiCl addition. (D-F) Skeletal muscle differentiation was rescued when the Wnt pathway was restored by LiCl in PCI treated zebrafish embryos. (D) Sarcomeric myosins, analysed by Western blot techniques, decreased in PCI treated embryos and returned comparable to those treated with DMSO when Wnt pathway was rescued adding LiCl (quantification in E). The efficacy of LiCl treatment was verified measuring the active β -catenin by Western blot techniques (quantification in F). Asterisks represent $*p < 0.05$, Student's t test.